*Fluid Phase Equilibria*, 94 (1994) 323–324
Elsevier Science B.V.



# Contents of Volume 94 (1994)

Phase equilibria of binary Lennard-Jones mixtures with cubic equations of state
V.I. Harismiadis (Amsterdam, The Netherlands), A.Z. Panagiotopoulos (Ithaca, NY, USA) and D.P. Tassios (Zographos, Greece)..... 1–18

Phase equilibria and thermodynamic properties of molecular fluids from perturbation theory. I. Pure fluids
K. Shukla (Houston, TX, USA)..... 19–49

Extension of Pitzer CSP models for vapor pressures and heats of vaporization to long-chain hydrocarbons
D.L. Morgan and R. Kobayashi (Houston, TX, USA)..... 51–87

A $g^E$ model for single and mixed solvent electrolyte systems. 1. Model and results for strong electrolytes
J. Li (Peking, China), H.-M. Polka and J. Gmehling (Oldenburg, Germany)..... 89–114

A $g^E$ model for single and mixed solvent electrolyte systems. 2. Results and comparison with other models
H.-M. Polka (Oldenburg, Germany), J. Li (Peking, China) and J. Gmehling (Oldenburg, Germany)..... 115–127

Solvation model for VLE in the system $H_2O$–LiBr from 5 to 76 wt%
R. Peters and J.U. Keller (Siegen, Germany)..... 129–147

Correlation of liquid–liquid phase equilibria using the SAFT equation of state
M.-L. Yu and Y.-P. Chen (Taipei, Taiwan)..... 149–165

Solid–liquid equilibria using DISQUAC. Prediction for 1-alkanol + *n*-alkane systems
J.A. González, I. García de la Fuente, J.C. Cobos, C. Casanova (Valladolid, Spain) and U. Domańska (Warsaw, Poland)..... 167–179

Prediction of VL and VLL equilibria of mixtures containing petroleum reservoir fluids and methanol with a cubic EoS
D. Avlonitis, A. Danesh and A.C. Todd (Edinburgh, UK)..... 181–216

Excess enthalpies of (ethanol or 1-propanol) + methyl *tert*-butyl ether + *n*-octane ternary mixtures at 298.15 K
S. Zhu, S. Shen, G.C. Benson and B.C.-Y. Lu (Ottawa, Ont., Canada) 217–226

Use of carbon dioxide to shift benzene/acetonitrile and benzene/cyclohexane azeotropes
P.G. Bendale and R.M. Enick (Pittsburgh, PA, USA)..... 227–253

Solubility and liquid density behavior for two binary systems of 2,2,4-trimethylpentane with methanol or nitroethane
H. Yoshikawa, T. Kanahira and M. Kato (Fukushima, Japan)..... 255–265

Liquid–liquid equilibria of ternary systems of water + acetone and $C_5$ and $C_8$ alcohols at 298, 303 and 308 K
A. Tiryaki, G. Gürüz and H. Orbey (Ankara, Turkey)..... 267–280

A new vapor–liquid equilibrium cell and VLE data for mixtures of 1-propanol + *p*-xylene
A.-Q. Chen, G.J. Urbanus and K.-C. Chao (West Lafayette, IN, USA). 281–288

Densities and excess volumes of the benzene–cyclohexane system between 298.15 and 473.15 K
S.A. Beg, N.M. Turkur, D.K. Al-Harbi and E. Hamad (Dhahran, Saudi Arabia) ..... 289–300

Calculation of helium solubilities using Henry's constants from cubic equations of state with generalized correlations for the interaction parameters
W. Schulze (Braunschweig, Germany) ..... 301–312

Liquid–liquid equilibria of the ternary system water/2-butyloxyethanol/*n*-decane at three different temperatures
S. Brandani, V. Brandani, G. Del Re and G. Di Giacomo (L'Aquila, Italy) ..... 313–320

*Subject Index* ..... 321

*Author Index* ..... 322

*Contents of Volume 94 (1994)* ..... 323–324